IMMUNIZATION REQUIREMENTS

Applicants enrolled or attempting to enroll shall have received the following vaccines in accordance with the doses and age requirements listed below. If, at any time, the age of the child is between the listed ages, the child must have received the number of doses in the "Total Doses Required" column.

| Institution | Age | Vaccine | Total Doses Required Column. | |
|--|--|--|--|--|
| | Less than 4 | Less than 4 This is not a recommended administration schedule, but contains the minimum requirements for participation in licensed child care. Routine vaccination | | |
| | months of age | begins at 2 months of age. | | |
| e Center | | Diphtheria/Tetanus/Pertussis | 1 dose | |
| | 4 months through 5 months of age | Polio | 1 dose | |
| | | haemophilus influenzae type B | 1 dose | |
| | | Pneumococcal | 1 dose | |
| | 6 months through 11 months of age | Dishtheric/Tetanya/Destyresia | 2 daga | |
| | | Diphtheria/Tetanus/Pertussis Polio | 2 doses 2 doses | |
| | | haemophilus influenzae type B | 2 doses | |
| | | Pneumococcal | 2 doses | |
| | | | | |
| | 12 months through 18 months of age | Diphtheria/Tetanus/Pertussis Polio | 3 doses 2 doses | |
| | | | 2 doses; or | |
| | | haemophilus influenzae type B | 1 dose received when the applicant is 15 months of age or older. | |
| | | | 3 doses if the applicant received 1 or 2 doses before 12 months of age; or | |
| Ca | | Pneumococcal | 2 doses if the applicant has not received any previous doses or has received 1 dose on or after 12 months of age. | |
| | | Diphtheria/Tetanus/Pertussis | 4 doses | |
| | 19 months through 23 months of age | Polio | 3 doses | |
| 7 | | | 3 doses, with the final dose in the series received on or after 12 months of age, or 1 dose received when the applicant is 15 | |
| Licensed Child Care | | haemophilus influenzae type B | months of age or older. | |
| | | | 4 doses; or | |
| | | Pneumococcal | 3 doses if the applicant received 1 or 2 doses before 12 months of age; or | |
| | | | 2 doses if the applicant has not received any previous doses or has received 1 dose on or after 12 months of age. | |
| | | Measles/Rubella ¹ | 1 dose of measles/rubella-containing vaccine received on or after 12 months of age; or the applicant demonstrates a | |
| | | | positive antibody test for measles and rubella from a U.S. laboratory. 1 dose received on or after 12 months of age if the applicant was born on or after September 15, 1997, unless the applicant | |
| | | Varicella | has had a reliable history of natural disease. | |
| | | | | |
| | 24 months and older | Diphtheria/Tetanus/Pertussis Polio | 4 doses | |
| | | | 3 doses 3 doses, with the final dose in the series received on or after 12 months of age; or 1 dose received when the applicant is 15 | |
| | | haemophilus influenzae type B | months of age or older. Hib vaccine is not indicated for persons 60 months of age or older. | |
| | | | 4 doses if the applicant received 3 doses before 12 months of age; or | |
| | | | 3 doses if the applicant received 2 doses before 12 months of age; or | |
| | | Pneumococcal | 2 doses if the applicant received 1 dose before 12 months of age or received 1 dose between 12 and 23 months of age; or | |
| | | | 1 dose if no doses had been received prior to 24 months of age. Pneumococcal vaccine is not indicated for persons 60 months of age or older. | |
| | | | 1 dose of measles/rubella-containing vaccine received on or after 12 months of age; or the applicant demonstrates a | |
| | | Measles/Rubella ¹ | positive antibody test for measles and rubella from a U.S. laboratory. | |
| | | Varicella | 1 dose received on or after 12 months of age if the applicant was born on or after September 15, 1997, unless the applicant | |
| | | varicella | has had a reliable history of natural disease. | |
| | | | | |
| Elementary or Secondary School (K-12) | | | 3 doses, with at least 1 dose of diphtheria/tetanus/pertussis-containing vaccine received on or after 4 years of age if the | |
| | | | applicant was born on or before September 15, 2000 ² ; or | |
| | 4 years of age and older | Diphtheria/Tetanus/ | 4 doses, with at least 1 dose of diphtheria/tetanus/pertussis-containing vaccine received on or after 4 years of age if the applicant was born after September 15, 2000, but before September 15, 2003 ² ; or | |
| | | Pertussis ^{4, 5} | applicant was born after September 15, 2000, but before September 15, 20032; or 5 doses with at least 1 dose of diphtheria/tetanus/pertussis-containing vaccine received on or after 4 years of age if the | |
| | | 1 GIUSSIS | applicant was born on or after September 15, 2003 ^{2, 3} ; and | |
| | | | 1 time dose of tetanus/ diphtheria/acellular pertussis-containing vaccine (Tdap) for applicants in grades 7 and above, if born | |
| | | | on or after September 15, 2000; regardless of the interval since the last tetanus/diphtheria containing vaccine. | |
| | | | 3 doses, with at least 1 dose received on or after 4 years of age if the applicant was born on or before September 15, 2003; | |
| | | Polio ⁷ | or | |
| | | | 4 doses, with at least 1 dose received on or after 4 years of age if the applicant was born after September 15, 2003.6 | |
| | | | 2 doses of measles/rubella-containing vaccine; the first dose shall have been received on or after 12 months of age; the | |
| | | Measles/Rubella ¹ | second dose shall have been received no less than 28 days after the first dose; or the applicant demonstrates a positive | |
| | | Hanatitis R | antibody test for measles and rubella from a U.S. laboratory. | |
| | | Hepatitis B | 3 doses if the applicant was born on or after July 1, 1994. 1 dose received on or after 12 months of age if the applicant was born on or after September 15, 1997, but born before | |
| | | | September 15, 2003, unless the applicant has had a reliable history of natural disease; or | |
| | | Varicella | 2 doses received on or after 12 months of age if the applicant was born on or after September 15, 2003, unless the | |
| | | | applicant has a reliable history of natural disease.8 | |
| Mumps vaccine | | easles/rubella-containing vaccine | · · · · | |

- 1 Mumps vaccine may be included in measles/rubella-containing vaccine.
- ² DTaP is not indicated for persons 7 years of age or older, therefore, a tetanus-and diphtheria-containing vaccine should be used.
- ³ The 5th dose of DTaP is not necessary if the 4th dose was administered on or after 4 years of age.
- 4 Applicants 7 through 18 years of age who received their 1st dose of diphtheria/tetanus/pertussis-containing vaccine before 12 months of age should receive a total of 4 doses, with one of those doses administered on or after 4 years of age
- 5 Applicants 7 through 18 years of age who received their 1st dose of diphtheria/tetanus/pertussis-containing vaccine at 12 months of age or older should receive a total of 3 doses, with one of those doses administered on or after 4
- 6 If an applicant received an all-inactivated poliovirus (IPV) or all-oral poliovirus (OPV) series, a 4th dose is not necessary if the 3th dose was administered on or after 4 years of age.
- ⁷ If both OPV and IPV were administered as part of the series, a total of 4 doses are required, regardless of the applicant's current age.
- Administer 2 doses of varicella vaccine, at least 3 months apart, to applicants less than 13 years of age. Do not repeat the 2nd dose if administered 28 days or greater from the 1nd dose. Administer 2 doses of varicella vaccine to applicants 13 years of age or older at least 4 weeks apart. The minimum interval between the 1nd dose of varicella for an applicant 13 years of age or older is 28 days.

 Revised 1-2013